

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



F.S.A.

FLUTRIAFOL 500 SC

FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL

GROUP **3** FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use

IMPORTANT: READ THIS BOOKLET BEFORE USING THIS PRODUCT



DIRECTIONS FOR USE: ALL STATES

F.S.A. FLUTRIAFOL 500 SC FUNGICIDE APPLIED AS A FERTILISER TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS												
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200 mL	4 weeks Grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of Granular Fertiliser (mL/100 kg)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333*</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL/100 kg)	60	333*	70	285	80	250	100	200	120	166
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL/100 kg)															
60	333*															
70	285															
80	250															
100	200															
120	166															

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS																				
Wheat	Stripe rust (<i>Puccinia striiformis</i> f.sp. <i>tritici</i>)	100 to 200 mL	4 weeks Grazing	<p>For control of strip rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.</p> <table border="1"> <thead> <tr> <th rowspan="2">Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100 kg of Granular Fertiliser (mL)</th> </tr> <tr> <th>100 mL rate</th> <th>200 mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>200</td> <td>400</td> </tr> <tr> <td>60</td> <td>166</td> <td>333</td> </tr> <tr> <td>70</td> <td>143</td> <td>285</td> </tr> <tr> <td>80</td> <td>125†</td> <td>250</td> </tr> <tr> <td>100</td> <td>100†</td> <td>200†</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL)		100 mL rate	200 mL rate	50	200	400	60	166	333	70	143	285	80	125†	250	100	100†	200†
	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL)																						
100 mL rate		200 mL rate																						
50	200	400																						
60	166	333																						
70	143	285																						
80	125†	250																						
100	100†	200†																						
	Septoria tritici blotch (<i>Mycosphaerella graminicola</i>) Take-all (<i>Geaumannomyces graminis</i>)	200 mL																						
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	100 to 200 mL		<p>* Application volumes over 300mL per 100kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.</p> <p>† Where application volume is low, product may be diluted with water to improve coverage.</p>																				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING

GENERAL INSTRUCTIONS

Fertiliser Application

Granulated Fertiliser

To determine the correct volume of F.S.A Flutriafol 500 SC Fungicide required to be applied, refer to Critical Comments in the Directions for Use table, which provides a guide based on a range of common fertiliser application rates. Ensure that treated fertiliser is dry before sowing with the crop.

Treated fertiliser should be applied in a band 3-4cm below or to the side of seed in planting furrow. It should not be broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertilizer and Flutriafol 500 SC.

Compatibility

Flutriafol 500 SC is compatible with the liquid fertilizers – Flexi N, Flexi NK, Flexi NS, Easy N and UAN. Agitation of the liquid fertilizer prior to and during addition of Flutriafol 500 SC is required. Maintain agitation of the mix during application.

Re-entry or Re-handling

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	----------	-----------

F.S.A. Flutriafol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management F.S.A Flutriafol 500 SC Fungicide is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to F.S.A Flutriafol 500 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by F.S.A. Flutriafol 500 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Four Seasons Agribusiness Pty Ltd accepts no liability for any losses that may result from the failure of F.S.A Flutriafol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

10L, 20L

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm

below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

110-1000L

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Four Seasons Agribusiness Pty Ltd should be advised immediately. This container is reusable and remains the property of Four Seasons Agribusiness Pty Ltd. DO NOT rinse empty container.

Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it.

When empty return to Four Seasons Agribusiness Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Harmful if swallowed. Harmful if inhaled. Avoid breathing fumes, mists, vapours or spray. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Not combustible. Use extinguishing media suited to burning materials. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used.

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com