

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



F.S.A.

ETHEPHON 900

GROWTH REGULATOR

ACTIVE CONSTITUENT: 900 g/L ETHEPHON
(an anti-cholinesterase compound)

For crop thinning, loosening or ripening in various crops and for accelerating boll opening, defoliation and pre-conditioning before defoliation of cotton as specified in the Directions for Use table.

IMPORTANT: READ THIS BOOKLET
BEFORE USING THIS PRODUCT



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to weak or stressed plants.

DO NOT apply F.S.A. Ethephon 900 following application of sodium chlorate, magnesium chlorate or diquat.

DO NOT apply to cotton if bolls have been killed by frost.

DO NOT apply if rain is expected within 8 hours of application.

DO NOT apply if day/night temperature is expected to be below 18°C.

DO NOT mix with hard (alkaline) water.

DO NOT apply before sufficient mature, unopened bolls have developed to produce the desired yield.

DO NOT pretreat with a desiccant (chlorate) before using F.S.A Ethephon 900 Growth Regulator.

CROP	ACTION	STATE	RATE (High Volume)	WHP	CRITICAL COMMENTS
APPLES: Jonathan, Delicious	Advancement of maturity, Improvement in red colour	Qld, NSW, Vic, SA, WA only	28 - 52 mL/100 L PLUS a non-ionic wetter and add "stop-drop"	7 days	Ensure fruit has reached marketable size before treatment and keep trees under observation after spraying. Apply 2 to 3 weeks before normal harvest period and 7 to 14 days before desired harvest date. Ensure fruit is picked at the correct stage of maturity, as fruit not harvested will quickly become over-ripe. NOTE: for best results spray when temperatures are between 15° and 32°C. Continuing low temperatures will advance maturity but may reduce colouration, so use lower rate. Use higher rate for quicker response. Add a "stopdrop" to prevent premature loosening of fruit. Thorough coverage is essential. Apply to fruit only for the fresh-fruit market or short-term storage.
Tydemán's Early		Tas only	20 mL per 100 L water and add "stop drop"		
Gravenstein, Golden Delicious, Jonathan and Red Delicious	Thinning*	Vic, SA only	12 mL/100 L water PLUS a non-ionic wetter		Apply at full bloom OR up to 1 week following full bloom. A second thinning spray 10 to 14 days after the F.S.A. Ethephon 900 spray may be necessary. On Gravenstein and Golden Delicious apply Carbaryl or 5 - 7.5 ppm NAA (use higher rate for maximum thinning). On Jonathan and Red Delicious apply Carbaryl or 4ppm NAA.
Golden Delicious, Stark Crimson and Leguna		Tas only	28 mL/100 L water a non-ionic wetter		Apply at the balloon blossom stage using a high volume sprayer at 4000 L/ha. DO NOT apply if rain is imminent. DO NOT use in sequence or in mixtures with NAA. If maximum temperatures are below 13°C after spraying, inadequate thinning may result, or if above 18°C excessive thinning may result.
Golden Delicious, Red Delicious, Jonathan, Granny Smith		WA only	28 mL to 32 mL/ 100 L water plus a non-ionic wetter		Only use high concentration when heavy set expected. * NOTE: for all thinning applications: The complex nature of fruit set and the possible variable action of chemical thinners due to weather conditions, make it difficult to give set recommendations and growers should consult their local agronomist.
Lady William			36 to 52 mL/100 L water plus a non-ionic wetter		

CROP	ACTION	STATE	RATE (High Volume)	WHP	CRITICAL COMMENTS
All varieties	Aid complete removal of fruit from trees and encourage a biennial bearing habit (Excessive vegetative growth will be suppressed and the following bloom stimulated).	NSW, SA, WA only	52 to 108 mL/ 100 L water PLUS a nonionic wetter	7 days	Apply just prior to full bloom or 5 to 6 weeks later during early morning or late afternoon, when slow drying conditions occur. Ensure thorough coverage of foliage and flowers or fruit. NOTE: Under normal weather conditions use lower rate for Delicious and higher rate for Jonathans as they are difficult to de-fruit. Consult the local Department of Agriculture for specific recommendations on other varieties. Warm weather following treatment will improve results, but if temperatures below 18°C prevail after spraying, fruit removal may be reduced; in the latter case use a higher dose rate. Fruit will loosen in 7 to 10 days and thinning should be complete within 2 to 3 weeks. Limbs can be lightly shaken to aid removal after this time.
	Retard vegetative growth and stimulate flowering of young apple trees in the following season				Apply from full bloom to 6 weeks after full bloom. Ensure thorough coverage. Trees should be large enough to support an increased crop of apples before being treated. Increase rate towards 108 mL/100 L to maximise thinning and stimulation of blooms during the following season. NOTE: Better results will be achieved if Daminozide at 1000ppm is added. DO NOT apply if harvesting fruit in the same season, as flowers or fruitlets will be partially or completely thinned.
CHERRIES: Ron's Seedling, St Margaret, Napoleon, Florence	Promotes evenness of maturity; early colour development	NSW only: Young district	16 mL/100 L water	7 days	Treatment with F.S.A. Ethephon 900: Apply when approximately 20% of the fruit show pink to red colour development; normally 7 to 10 days before harvest. Ensure thorough coverage of leaves and fruit. NOTE: high temperature at spraying or following spraying may accelerate results. DO NOT spray when mean daily temperatures exceed 32°C. DO NOT spray when rain is expected within 24 hours OR when temperatures are low (17°C or less) as this will delay or reduce the effect of F.S.A. Ethephon 900. Fruit will mature 3 to 5 days earlier than normal and will be loosened to allow easier harvesting. A high proportion of fruit will readily separate from the stalk when harvested.
		Orange/Bathurst district	32 mL/100 L water		
TABLE GRAPES: Barlinka	To promote early, uniform colouring	NSW, SA only	12 to 16 mL per 100 L water PLUS a non-ionic wetter	14 days	Apply when 5 to 30% of the berries are coloured (2 - 4 weeks prior to expected harvest). Use the higher rate to maximise the colouring effects required. Ensure thorough coverage of bunches although the foliage need not be completely covered. NOTE: F.S.A. Ethephon 900 may slightly reduce firmness of grapes. DO NOT use on grapes intended for long storage.
Red Emperor, Red Prince, Red Malaga		NSW, Vic, SA, WA only	20 to 32 mL/ 100 L water PLUS a non-ionic wetter		
Flame Seed-less		WA only			

CROP	ACTION	STATE	RATE (High Volume)	WHP	CRITICAL COMMENTS
WINE GRAPES: Semillon	Aids to mechanical harvesting	NSW only	32 to 104 mL PLUS a non-ionic wetter	7 days	Apply 7 days before expected harvest. Thorough coverage of upper and lower leaf surfaces is essential. NOTE: some leaf yellowing and leaf fall, which is not detrimental to vines, may occur. The concentration to use will depend on a number of factors such as seasonal conditions, crop size, weather conditions before and after application, and trellis type. Therefore, before applying consult the local Department of Agriculture for specific recommendations.
MANDARINS Imperial	Thinning to increase fruit size, to reduce size of heavy crop and to even out the production cycle	Qld, NSW, SA, WA only	28 to 32mL/ 100 L water	Not re-quired	Apply during the early stage of fruit development for thinning and correction of the habit of alternate heavy and light crops. Apply as foliar spray when fruitlets are about 10 to 15 mm in diameter and when natural fruit drop is occurring. This is usually in November in Qld, December in SA and intermediate in NSW, WA and Vic. Weather conditions influence the degree of thinning and care must be taken not to over-thin. NOTE: Fruitlets should fall off 7 to 14 days after application. Aim to apply 13 to 15 L of spray per 4 to 5 m high tree. Use the higher rate when a very heavy crop is evident. DO NOT apply in cold weather (less than 18°C) OR when rain is likely within 1 to 2 days of spraying. DO NOT add surfactants or wetting agents as leaf fall may be increased. See also General Instructions.
		Vic only	20 to 28 mL/ 100 L water		
ORANGES Navel		NSW, WA only	28 to 32 mL/ 100 L water		
Valancia		NSW, SA, WA only			
		Vic only	32 to 36 mL/ 100 L water		
MACADAMIA NUTS: Own choice	Aid harvesting by promoting uniform nutfall	NSW, WA only	132 mL/100 L water PLUS a non-ionic wetter	7 days	Apply late March to early May when nuts are mature. Vary rate according to degree of loosening required. NOTE: Nuts will be stimulated to fall within 10 to 14 days after spraying. Mechanical shaking may be used 7 to 10 days after spraying. Ensure thorough coverage of foliage and nuts. Ensure tops and insides of trees are adequately sprayed. A small quantity of older leaves will fall after treatment, but production will not be adversely affected. DO NOT use on Teddington variety.
H2 Variety			44 to 84 mL/ 100 L water PLUS a non-ionic wetter		
PEACHES	Advancement and concentration of maturity	VIC only: Goulburn Valley	12 mL/ 100 L water PLUS a non-ionic wetter	6 weeks	Apply once as foliar spray after commencement of the final fast growth stage. Timing depends on variety. Determine by measuring twice weekly, fruit circumference, of 20 tagged fruit/block. After rapid growth stage determined, wait 3 to 4 days for further confirmation, then spray. Thorough coverage is essential. NOTE: Instances have been recorded where this product, when applied in peaches at apparently the correct time, has resulted in premature fruit drop, fruit gumming and fruit splitting. See also General Instructions.

CROP/ SITUATION	ACTION	STATE	RATE		WHP	CRITICAL COMMENTS
			General	Boom Spraying		
Pineapples (plus Urea)	Initiation of flowering	Qld, WA only	64 mL + 5kg Urea /100 L water Apply 30 mL solution/plant	1.28 L PLUS 100 kg Urea in 2000 L of water/ ha (Boom spray)	7 days	Apply in March. Note: Rate should be doubled for plant crops that are growing vigorously at the time of application.
			32 mL + 5kg Urea /100 L water Apply 30 mL solution/plant	640 mL PLUS 100 kg Urea in 2000 L of water/ ha (Boom spray)		Apply in May - June OR September - October. Note: See aboe - February/March application with Urea
Pineapples (without urea)	Initiation of flowering in ratoon	Qld, WA only	20 mL/10 L water Apply 30 mL solution/plant	4.8 L in 2000 L of water/ha (Boom spray)	7 days	Apply February – March. Note: Use on the final crop in the block before eradication of the crop. Pineapples treated at the above time will be ready for picking 7 to 10 months after application.
			6 mL/10 L water Apply 30 mL solution/plant	1.2 L in 2000 L of water/ha (Boom spray)		Apply May – June OR September – October. Note: See above – February/March application without Urea.
Pineapples	Fruit ripening		-	1.36 L in 1000 L water/ha		F.S.A. Ethephon 900 for ripening should only be used on even crops successfully induced for flowering with F.S.A. Ethephon 900. DO NOT use on fields intended for ratoon production. Treatment should be made when the forced fruit are beginning to show the first colour break. An initial harvest may be necessary to remove more advanced fruit before applying F.S.A. Ethephon 900. The remaining treated fruit will be ready for harvest at least 7 days following the F.S.A. Ethephon 900 ripening application. Note: DO NOT use on fruit intended for the fresh fruit market. F.S.A. Ethephon 900 for ripening is NOT SUITABLE FOR USE ON SUMMER PLANT CROPS because of scattered fruiting and high risk of forcing young suckers. Where natural fruiting has occurred prior to chemical induction, these natural fruit must be picked before F.S.A. Ethephon 900 ripening. Treatment too early can result in a loss in yield and a reduction in fruit sugar levels.

CROP/ SITUATION	USE	STATE	RATE (High Volume)	WHP	CRITICAL COMMENTS
Sugarcane	Cane ripening	Qld, NSW, WA only	800 mL/ha	6 weeks	Use in March-April to accelerate ripening in cane and increase c.c.s. Use on cane planted for early harvest with harvest no later than August. Apply to actively growing and non-stressed cane with at least eight non-stressed green leaves. Cane varieties Q115, Q119, Q137, Q124, Q96, H56-752 and CP44-101 have all shown high to moderate response to ethephon. Shortening of some internodes and yellowing of leaves may occur without any effect on cane yield. Test any new varieties prior to application.
Tomatoes	Accelerate ripening, increasing yield of marketable fruit	All States	960 mL in 440 to 880 L water/ha	7 days	Apply once in the season. Ensure thorough coverage of fruit and foliage, when fruit is 5 to 30% pink or red. To determine this, pull a few plants, shake off all fruit, weigh and determine above percentage. Optimum harvest maturity is expected 14 to 21 days after spraying. DO NOT use on greenhouse crops. Note: Temperatures below 18°C retard colour development and may extend the interval between treatment and harvest. Treatment may cause some yellowing of foliage or defoliation. Sun scald or exposed fruit may occur under high temperature conditions following treatment.

CROP/ SITUATION	USE	STATE	RATE (High Volume)	CRITICAL COMMENTS
Cotton	Defoliation, acceleration of boll opening	Qld, NSW, WA, NT only	1.6 – 2.4 L/ha in a minimum of 20 L of water/ha	FOR ENHANCED DEFOLIATION AND BOLL OPENING OF COTTON. Apply when the number of mature unopened bolls is sufficient to produce the desired crop. Use the lower rate in warm, optimum conditions. Use the higher rate for stressed plants, rank growth, or adverse cool conditions (eg if day/night temperature after treatment is expected to be below 22°C). Ensure thorough and uniform coverage of bolls.
	Pre-conditioning of crop before a normal defoliation is to occur		1.04 L/ha in a minimum of 20 L of water/ha	FOR PRE-CONDITIONING BEFORE DEFOLIATION. Apply when there are sufficient mature unopened bolls to produce the desired yield. Apply defoliant 4-7 days later. Ensure thorough and uniform coverage of bolls.
	Defoliation #		400 to 800 mL/ ha in a minimum of 20 L of water/ ha	Use in a tank mix with a registered rate of thidiazuron. Apply when there are sufficient unopened bolls to produce the desired yield. Total application volume should be no less than 20 L/ha. Ensure thorough and uniform coverage of bolls and leaves. # This mixture will also provide some acceleration of boll opening, however, a subsequent application of a boll opener may be required to complete boll opening.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

APPLES, CHERRIES, MACADAMIA NUTS, PINEAPPLES, TOMATOES, WINE GRAPES:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

TABLE GRAPES AND COTTON: **DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

PEACHES AND SUGAR CANE: **DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.**

GRAZING: **DO NOT FEED COTTON STUBBLE OR GIN TRASH TREATED WITH F.S.A. ETHEPHON 900 TO LIVESTOCK.**

DO NOT GRAZE LIVESTOCK ON TREATED AREAS.

GENERAL INSTRUCTIONS

When adding a non-ionic wetter, use the rate as recommended on the wetter label.

IMPORTANT

Apples- advancement of fruit maturity and improvement of red colour:

- Ensure fruit has reached marketable size before treatment and keep trees under observation after spraying.
- Ensure fruit is picked at the correct stage of maturity, as fruit not harvested will quickly become over-ripe.
- Continuing low temperatures will advance maturity and colouration; continuing warm to high temperatures, both day and night, will advance maturity but may reduce colouration.
- Apply to fruit only for the fresh fruit market or short-term storage.

Apples - Thinning:

The complex nature of fruit and the possible variable action of chemical thinners due to weather conditions, make it difficult to give set recommendations and growers should consult their local agronomist.

Apples - Fruit Removal:

- Apply to trees to remove unwanted fruit and encourage a biennial bearing habit. Excessive vegetative growth will be suppressed and bloom the following season stimulated.
- Weather conditions are very important – warm weather following treatment will improve results, but if temperatures below 18°C prevail after spraying, fruit removal may be reduced. In the latter case, use the higher dose rate.
- Fruit will loosen in 7-10 days and thinning should be complete within 2-3 weeks. Limbs can be lightly shaken to aid removal after this time.

Imperial Mandarins and Valencia Oranges:

Apply during the early stage of fruit development for thinning and correction of the habit of alternate heavy and light crops. Weather conditions influence the degree of thinning and care must be taken not to over-thin.

EXCLUSION OF LIABILITY:

As the degree of thinning may vary according to the conditions and over-thinning is possible, the decision to incur these risks is taken by the user, and the manufacturer and vendor exclude themselves from all conditions, representations, warranties, whether expressed or implied by the statute or otherwise and accept no responsibility for any losses which follow from the use of the product on mandarins or oranges as far as this is legally possible under the Trade Practices Act or any State legislation.

Cotton:

A foliar spray with F.S.A. Ethephon 900 will accelerate opening of mature unopened bolls and enhance defoliation, which can result in earlier harvest and increased recoverable yield and quality. Some premature drop of small immature bolls may occur.

Harvest at optimum boll opening and optimum defoliation.

Too late harvest may reduce quality and lint.

Cotton - APPLICATION

DO NOT apply before sufficient mature unopened bolls have developed to produce the desired yield of cotton. Bolls are mature when the seed coat turns light brown, the embryo fills the seed cavity with an abundance of creamy colour and cotyledons are well defined.

Cotton - Treatment sequence

- Light canopy: Apply F.S.A. Ethephon 900 then apply defoliant 4 to 7 days later.
- Moderate to heavy canopy: First apply a pre-conditioner, then apply F.S.A. Ethephon 900 as soon as sufficient leaves have dropped to expose the target bolls, usually 4 to 10 days later.

Cotton - APPLICATION METHOD

Apply by placement technique under the following range of conditions to achieve a target droplet density of 60 droplets per sq.cm.

Wind Direction: Cross Wind Wind speed range: 6-20kph

Air Temperature: Maximum 32°C Delta t: Maximum 10°C

Sugar Cane:

F.S.A. Ethephon 900 may be applied by air or ground spray equipment. Apply in 30 L water/ha by air, and 110-150 L water/ha by ground spray equipment.

Apply during March-April on cane due for harvest no later than August. Consult your local Four Seasons Agribusiness agent or BSES officer.

Tomatoes:

- Temperatures below 18°C retard colour development and may extend the interval between treatment and harvest.
- Treatment may cause some yellowing of foliage or defoliation.
- Sun-scald of exposed fruit may occur under high temperature conditions following treatment.

MIXING AND APPLICATION

Add F.S.A. Ethephon 900 to half the required volume of good quality water in the spray tank, then add the remaining volume of water while agitating the mixture. DO NOT use alkaline water. Prepare only enough spray solution for immediate use, do not allow to stand.

If tank mixes are prepared maintain good agitation and use promptly. Apply as a thorough uniform spray to achieve good coverage. For aerial application apply in at least 20 L/ha water. F.S.A. Ethephon 900 is corrosive to acrylic plastics, certain paints and metals. Flush and rinse all exposed equipment including aircraft surfaces with detergent and water after use.

COMPATIBILITY

DO NOT mix F.S.A. Ethephon 900 and sodium chlorate as this could result in emission of toxic chlorine fumes. F.S.A. Ethephon 900 may be tank-mixed with defoliant products. Since the formulations of other products are beyond the control of Four Seasons Agribusiness Pty Ltd tank mixtures should be tested before commercial use.

CAUTION

EXPORT OF TREATED PRODUCE

Growers should note the Maximum residue limits (MRL's) or import tolerances do not exist in all markets for products treated with F.S.A. Ethephon 900. Additionally, some export markets have established MRL's that are different to those in Australia.

If you are growing product for export (either fresh, dry or for wine production), please check with your Four Seasons Agribusiness agent or industry body for the latest information on MRL's and Import tolerances BEFORE using F.S.A. Ethephon 900.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store in direct sunlight.

Store in the closed container in a dry, cool, well ventilated area away from children, animal, food, feedstuffs, seed and fertilisers.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. Will damage eyes and skin. Repeated minor exposure may have a cumulative poisoning effect.

Do not inhale vapour spray mist. Avoid contact with eyes and skin. When opening the container and preparing spray and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and chemical resistant footwear and face shield or goggles. If product or spray in eyes, wash it out immediately with water. If product or spray on skin, immediately wash area with soap and water.

Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If swallowed, do NOT induce vomiting. If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised.

Give Atropine if instructed. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which is available from the supplier.

CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Causes severe skin burns and eye damage. Harmful if inhaled. Harmful to aquatic life with long lasting effects. Do not breathe fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Use only outdoors or in a well ventilated area. Avoid release to the environment. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com