

# POISON

KEEP OUT OF REACH OF CHILDREN

FOR ANIMAL TREATMENT ONLY

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# THREEXTRIPLE

## Combination Drench For Sheep

### ACTIVE CONSTITUENTS:

34 g/L Levamisole

(equivalent to 40 g/L Levamisole hydrochloride)

25 g/L Albendazole

1 g/L Abamectin

0.5 g/L Selenium (as Sodium Selenate)

0.21 g/L Cobalt (as Cobalt Sulphate Heptahydrate)

0.6 g/L Zinc (as Zinc Sulphate Monohydrate)

0.76 g/L Iodine (as EDDI stabilised)

For the control of major gastrointestinal parasites in sheep, including strains with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families, susceptible to the combination of Abamectin, Levamisole and Albendazole. Also controls lungworm, itch mite, larval stages of nasal bot. Provides a supplemental source of selenium, zinc, iodine and cobalt.



FOUR SEASONS  
AGRIBUSINESS

# THREE X TRIPLE

## Combination Drench For Sheep

Three XTriple Combination Drench for Sheep contains levamisole, albendazole and abamectin, which are members of the imidazothiazole, benzimidazole and macrocyclic lactone groups respectively.

For the effective control of mature and immature stages of susceptible strains of the following parasites

### Gastrointestinal Parasite:

Barber's pole worm *Haemonchus contortus* (Including inhibited L4 stage)  
Small brown stomach worm *Teladorsagia circumcincta* (Including inhibited L4 stage)  
Stomach hair worm *Trichostrongylus axei*  
Black scour worm *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*  
Thin necked intestinal worm *Nematodirus spathiger*, *Nematodirus filicollis*  
Small intestinal hair worm *Cooperia curticei*, *Cooperia oncophora*  
Intestinal threadworm *Strongyloides papillosus*  
Large mouthed bowel worm *Chabertia ovina*  
Nodule worm *Oesophagostomum columbianum*  
Whipworm *Trichuris ovis*  
Nasal bot *Oestrus ovis* (parasitic larval stages only)  
Large lungworm *Dictyocaulus filaria*  
Itchmite *Psorergates ovis*

Resistance may develop to any chemical.

Ask your local Veterinary practitioner or Animal Health Advisor for recommended management practices. Check the effectiveness of anthelmintic at least annually and it is advised that a drench resistance test be conducted before any drench is used.

### DIRECTIONS FOR USE:

#### Restraints:

**MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.**

**CONTRAINDICATIONS:** This product is contraindicated for use in lambs under 6 weeks of age or less than 15 kg body weight.

This product should not be used when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

This product is contraindicated for use unless cobalt deficiency is confirmed. This product is contraindicated for use in sheep that are dehydrated or unwell. This product is contraindicated for use unless mineral deficiency is confirmed. Dogs, horses or any other pets should not be drenched with this product.

This product is not recommended for use in goats as safety and efficacy have not been evaluated.

**PRECAUTIONS:** Recommended dose should not be exceeded.

Minerals such as cobalt, selenium, iron, manganese and molybdenum compete with each other for metabolically important processes in the animal. Users should seek veterinary advice before treatment.

Fatal interactions may occur between levamisole and organophosphate dips. Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.

Do not treat sheep at intervals of less than 3 weeks, due to selenium content.

### DOSAGE AND ADMINISTRATION:

#### SHAKE WELL BEFORE USE.

For oral administration only.

Dose rate is 1ml per 5kg bodyweight equivalent to 8mg Levamisole hydrochloride, 5mg Albendazole, 0.2mg abamectin per kg bodyweight.

Body Weight (kg)	Dose (mL)	Body Weight (kg)	Dose (mL)
15	3	46-50	10
16-20	4	51-55	11
21-25	5	56-60	12
26-30	6	61-65	13
31-35	7	66-70	14
36-40	8	71-75	15
41-45	9		

Over 75 kg add 1mL for every 5kg bodyweight

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs); (bucks, does, kids). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Do not re-treat animals for 28 days after last treatment.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS:

**MEAT: DO NOT USE less than 14 days before slaughter for human consumption.**

### TRADE ADVICE:

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. Before using this product confirm the current ESI from Four Seasons Agribusiness Pty Ltd on 1300 449 255 or the APVMA website [www.apvma.gov.au/residues](http://www.apvma.gov.au/residues).**

**SAFETY DIRECTIONS:** Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID INSTRUCTIONS:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**ADDITIONAL USER SAFETY:** Additional information is listed in Safety Data Sheet.

**ENVIRONMENTAL PROTECTION:** Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

**DISPOSAL:** Triple-rinse container into the medicated drench. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**STORAGE:** Store below 25°C (air conditioning). Do Not Freeze. Protect from Light.

**CONDITIONS OF SALE:** The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No.: 94927/144054

**IN AN EMERGENCY, DIAL 000, POLICE OR FIRE BRIGADE**



Four Seasons Agribusiness Pty Ltd  
287 Boorowa Street, Young, NSW, Australia, 2594  
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255  
[www.fourseasonsag.com](http://www.fourseasonsag.com)