

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Quad

Multi-Combination Sheep Drench

ACTIVE CONSTITUENTS: 37.5 g/L closantel
34 g/L levamisole (as levamisole hydrochloride)
25 g/L albendazole
1 g/L abamectin

For the treatment and control in sheep of susceptible gastrointestinal roundworms (including strains with single or dual resistance to macrocyclic lactones, benzimidazoles, levamisole or closantel, and strains of *Haemonchus contortus* with emerging resistance to closantel). It is also effective against lungworm, tapeworms, mature and late immature liver fluke, nasal bot and itch mite.



10L



FOUR SEASONS
AGRIBUSINESS

QUAD MULTI-COMBINATION SHEEP DRENCH

QUAD Multi-Combination Sheep Drench contains abamectin, albendazole, closantel and levamisole, which are members of the macrocyclic lactone, benzimidazole, salicylanilide and imidazothiazole family of chemicals respectively.

It is a broad-spectrum oral anthelmintic which can be used against susceptible gastrointestinal parasites, and to prevent the introduction of resistant strains on to a property.

It is effective against sensitive strains of the following parasites of sheep:

Adult and immature gastrointestinal roundworms

- Barber's Pole Worm (*Haemonchus contortus*) - including inhibited L4 stage and strains with single, dual or triple resistance to macrocyclic lactones, benzimidazoles, levamisole or closantel, and strains with emerging resistance to closantel.
- Small Brown Stomach Worm (*Teladorsagia (Ostertagia) circumcincta*) - (including inhibited L4 stage and strains with single or dual resistance to macrocyclic lactones, benzimidazoles or levamisole.
- Black Scour Worm (*Trichostrongylus colubriformis*) including strains with single or dual resistance to macrocyclic lactones, benzimidazoles or levamisole.
- Large Stomach Worm (*Haemonchus placei*)
- Stomach Hair Worm (*Trichostrongylus axei*)
- Black Scour Worm (*Trichostrongylus vitrinus*)
- Small Intestinal Worm (*Cooperia* spp.)
- Thin-Necked Intestinal Worm (*Nematodirus* spp.)
- Large-mouthed bowel worm (*Chabertia ovina*)
- Nodule Worm (*Oesophagostomum columbianum*)
- Large Bowel Worm (*Oesophagostomum venulosum*)
- Whip Worm (*Trichuris ovis*)
- Intestinal threadworm (*Strongyloides papillosus*)
- Hookworm (*Bunostomum* spp.)

Nasal Bot (*Oestrus ovis*)

Lungworm - Large lungworm (*Dictyocaulus viviparus*)

Liver Fluke (*Fasciola hepatica*) - mature fluke and immature forms including 6 week old stages. Reduces the output of viable fluke eggs.

Tapeworms (*Moniezia* spp.) - heads and segments

Itch mite (*Psorergates ovis*)

Resistance may develop to any chemical. Ask your local veterinarian or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

DIRECTIONS FOR USE:

Restraints: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

CONTRAINDICATIONS: This product is contraindicated for use in lambs under 6 weeks of age or less than 10 kg body weight.

PRECAUTIONS: Use with caution and exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.

Avoid yarding animals off-feed overnight, and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded, except under veterinary supervision.

SHAKE WELL BEFORE USE

Dose: 1 mL per 5 kg bodyweight.

Administer orally.

Check the accuracy of the drench equipment before and during use.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range.

Do not underdose. Drafting into 2 or more lines may be appropriate, to avoid excessive overdosing.

DOSAGE TABLE

Liveweight	Dose	10L treats
11 - 15 kg	3 mL	3333
16 - 20 kg	4 mL	2500
21 - 25 kg	5 mL	2000
26 - 30 kg	6 mL	1666
31 - 35 kg	7 mL	1428
36 - 40 kg	8 mL	1250
41 - 45 kg	9 mL	1111
46 - 50 kg	10 mL	1000
51 - 55 kg	11 mL	909
56 - 60 kg	12 mL	833
61 - 65 kg	13 mL	769
66 - 70 kg	14 mL	714
71 - 75 kg	15 mL	666

Sheep in excess of 75 kg to be dosed at 1 mL per 5 kg bodyweight.

GENERAL DIRECTIONS:

For use as a quarantine drench: Treat all animals on arrival, or ideally at farm of origin, and hold in yards for 24 hours before release.

For use in a strategic worm control program: Speak to your animal health adviser or veterinarian.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product.

Note - observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Four Seasons Agribusiness Pty Ltd on 1800 560 708 before using this product.

SAFETY DIRECTIONS: POISONOUS IF ABSORBED BY SKIN

CONTACT OR SWALLOWED. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and pouring large quantities wear elbow-length PVC gloves. After each days use, wash gloves. Wash hands after use.

FIRST AID INSTRUCTIONS: If poisoning occurs, get to a doctor or hospital quickly.

ADDITIONAL USER SAFETY: CAN CAUSE PERMANENT VISION LOSS IF SWALLOWED.

Additional information is listed in the safety data sheet (SDS)

ENVIRONMENTAL PROTECTION: Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used containers.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Store below 30°C (room temperature), in original container. Do not freeze. Do not store at low temperature.

CONDITIONS OF SALE: The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No.: 95108/144699

Additional GHS Information

Harmful if inhaled. May damage fertility or the unborn child.

If medical advice is needed, have product container or label at hand. Avoid breathing vapours. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Store locked up.

TECHNICAL SUPPORT: 1800 560 708

**IN AN EMERGENCY, DIAL 000
POLICE OR FIRE BRIGADE**



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, NSW, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189
Phone 1300 449 255
www.fourseasonsag.com